arXiv:1304.2933v1 [cond-mat.str-el] 10 Apr 2013

Semi-metal-insulator transition on the surface of a topological insulator with in-plane magnetization

Flavio S. Nogueira and Ilya Eremin
Institut für Theoretische Physik III, Ruhr-Universität Bochum, Universitätsstraße 150, DE-44801 Bochum, Germany

(Dated: Received April 11, 2013)

A thin film of ferromagnetically ordered material proximate to the surface of a three-dimensional topological insulator explicitly breaks the time-reversal symmetry of the surface states. For an out-of-plane ferromagnetic order parameter on the surface, parity is also broken, since the Dirac fermions become massive. This leads in turn to the generation of a Chern-Simons term by quantum fluctuations. On the other hand, for an in-plane magnetization the surface states remain metallic. In this work we study the possibility of spontaneous breaking of parity due to a dynamical gap generation on the surface. We find that in the absence of interaction between the fermions there is no spontaneous gap generation. In the presence of a local, Hubbard-like, interaction of strength g, a gap and a Chern-Simons term are generated for g larger than some critical value, g_c, provided the number of Dirac fermions, N, is odd. For an even number of Dirac fermions the masses are generated in pairs having opposite signs, and no Chern-Simons term is generated. Our result offers a possible explanation to recent experiments showing a gap opening even when the topological insulator is proximate to a planar ferromagnet.

PACS numbers: 75.70.-i,73.43.Nq,64.70.Tg,75.30.Gw

Due to its unique properties, topological insulators (TI) [1, 2] are likely to play a major role as a component material in different types of heterostructures. For instance, with a view towards spintronics applications [3], heterostructures involving ferromagnetic (FM) materials or magnetic impurities have been studied both theoretically [4–13] and experimentally [14–20]. Assuming a strong spin-orbit coupling, the Lagrangian of the TI surface proximate to a FM thin film inducing a magnetization $\mathbf{n} = (n_x, n_y, \sigma)$ on it is given by,

$$\mathcal{L} = \bar{\psi}[i\gamma_0\partial_t - i\vec{\gamma}\cdot(v_F\nabla + iJ\mathbf{a}) - J\sigma]\psi, \quad (1)$$

where $\psi = [c_\uparrow, c_\downarrow]^T$ and $\bar{\psi} = \psi^\dagger\gamma^0$, with $\gamma^0 = \sigma_z$, $\gamma^1 = -i\sigma_x$, and $\gamma^2 = i\sigma_y$. The above Lagrangian has a QED-like form in $d = 2 + 1$ dimension with a vector potential $\mathbf{a} = (n_y, -n_x)$, and no time component for the gauge field. The magnetization $\mathbf{n}(t, \mathbf{r})$ originates from a proximity effect with a FM thin film over the TI surface. Note that time-reversal symmetry is only broken if $\sigma \neq 0$. Thus, an induced planar magnetization does not break time-reversal invariance.

If the electrons on the surface of the topological in are assumed to interact via a local, Hubbard-like repulsive interaction,

$$\mathcal{L}_{\rm int} = -\frac{g}{2}(\bar{\psi}\gamma^0\psi)^2, \quad (2)$$

where $g > 0$, we can introduce an auxiliary field, a_0, via a Hubbard-Stratonovich transformation to obtain,

$$\mathcal{L} = \bar{\psi}[i\gamma^0(\partial_t + iJa_0) - i\vec{\gamma}\cdot(v_F\nabla + iJ\mathbf{a}) - J\sigma]\psi - \frac{J^2}{2g}a_0^2. \quad (3)$$

Now the electronic sector is fully gauge-invariant. The only term in the Lagrangian that (explicitly) breaks the gauge symmetry is the Gaussian factor in the auxiliary field a_0. One consequence of this term, quadratic in a_0, is that g does not renormalize. This is a consequence of the gauge symmetry of the fermionic sector and can be easily proved using Ward identities [21]. An easy way of seeing this without making explicit use of the Ward identity is to introduce renormalized fields $\psi_r = Z^{-1/2}\psi$ and $a_r^\mu = Z_a^{-1/2}a^\mu$ and observe that gauge invariance of the fermionic sector implies that the renormalized exchange coupling is $J_r = \sqrt{Z_a}J$. This implies at the same time that terms quadratic in a^μ which are not gauge invariant should not renormalize, yielding $J_r^2/g_r = Z_aJ^2/g$ and consequently $g_r = g$. Therefore, g is a good tuning parameter in our theory. The fact that g does not renormalize will be important in our subsequent analysis.

Note that we are disregarding the long-range contribution to the Coulomb interaction. Such a long-range contribution has been shown to be irrelevant in the long wavelength limit in theoretical studies of interacting graphene [22–24] and a similar reasoning also applies here. Our Lagrangian corresponds to a restricted Thirring model [25], in the sense that only the zeroth component of the current $j_\mu = \bar{\psi}\gamma_\mu\psi$ appears squared in the interaction.

From the Lagrangian (3) it is readily seen that in the case of an out-of-plane magnetization ($\sigma \neq 0$), the Dirac fermions become massive when the symmetry is spontaneously broken, in which case the expectation value $\langle\sigma\rangle \neq 0$. On the other hand, in-plane magnetization ($\sigma = 0$) implies massless Dirac fermions; see Fig. 1. An interesting question to ask is whether a mass can also be generated in this case. The answer to this question is far less obvious than in the case of out-of-plane magnetization. This point relates to similar studies in (2+1)-dimensional, massless, QED-like theories in diverse condensed matter physics contexts, ranging from high-T_c cuprates [7, 26, 27, 29] to interacting graphene [30–33]. However, in the graphene case only the zeroth component of the gauge field is present.

The case of in-plane magnetization is of great experimental relevance. Recently a thin film of FM insulator EuS has been successfully grown on the surface of $\mathrm{Bi_2Se_3}$ [20], making the surface of $\mathrm{Bi_2Se_3}$ ferromagnetic by proximity effect. The high magnetic anisotropy of the EuS thin film essentially induces an in-plane magnetization on the topological surface.



(a) Out-of-plane magnetization: gapped Dirac spectrum



(b) In-plane magnetization: gapless Dirac spectrum

FIG. 1. (Color online) Schematic comparison between two types of magnetization orientation on the surface of a TI. (a) In the case of out-plane magnetization the electronic spectrum at the surface is gapped. (b) For in-plane magnetization the spectrum is gapless.

In spite of this high magnetic anisotropy, the experiments of Wei *et al.* [20] find a large gap in the Dirac spectrum. In this paper we show that such a gap may be spontaneously generated in FM/TI heterostructures by a parity symmetry breaking mechanism if interaction effects are taken into account. Our analysis shows that the long sought evidence for such spontaneous gap generation in QED-like theories may be within experimental reach in TIs.

QED-like theories in two spatial dimensions feature either two- or four-component Dirac fermions. Massless QED in 2+1 dimensions with four-component Dirac fermions has a chiral symmetry, which prevents the addition of a mass term [37]. Indeed, a term $m\bar{\psi}\psi$ in the Lagrangian would explicitly break the chiral symmetry [37]. However, it can be shown that a fermion mass is dynamically generated, breaking in this way the chiral symmetry spontaneously. This occurs if the number of Dirac fermion species, N, is smaller than some critical value [36]. In the context of high-T_c superconductors this mass generation mechanism is associated to the development of antiferromagnetic order [26, 28, 29, 34]. In the graphene case it is associated to an emergent excitonic gap [24, 30–32].

The QED-like theory we consider here features two-component Dirac fermions and, for this reason, does not have a chiral symmetry [36, 37], as γ_5-like matrices can only be defined in representations featuring four-component spinors [21] (see, however, the discussion below on the number of Dirac fermion species). Hence, the massless case corresponding to an in-plane magnetization ($\sigma = 0$) has no more *internal* symmetry than the massive case corresponding to an out-of-plane magnetization. On the other hand, a mass term breaks the parity symmetry, such that integrating out the fermions generates a Chern-Simons (CS) term [38], inducing in this way a topological magnetoelectric effect [4]. This is what happens in the case of an out-of-plane magnetization, where the generation of the CS term significantly affects the magnetization dynamics [11]. Therefore, it seems that a dynamical mass generation in the case of in-plane magnetization would spontaneously break the parity symmetry. For the case of a theory containing N two-component massless Dirac fermions, it was found in the past using a large N approach that masses are indeed dynamically generated, but half of them acquire a mass $+m$ while the other half acquire a mass $-m$, implying that the parity symmetry is actually not broken [39]. A recent more refined study at finite N [40] confirms this result for $N_+ = N_-$, where $N_\pm$ denote the number of fermions with mass $\pm m$, respectively. As a consequence, no CS term is generated, since the CS mass is proportional to $\sum_{a=1}^{N} m_a/|m_a|$ and have no further corrections arising from quantum fluctuations, as required by the Coleman-Hill theorem [41]. It is also possible to assume that some kind of chiral symmetry is broken in the case of two-component Dirac fermions by considering instead a representation featuring $N/2$ four-component Dirac fermions in a reflection-invariant vacuum state [39, 40]. The numerical solutions of the Schwinger-Dyson equation [40] in this case yield parity violation when $(N_+ - N_-)/(N_+ + N_-) < 0.4$. However, this result seems to be in disagreement with a theorem by Vafa and Witten [42], which would exclude spontaneous parity symmetry breaking in gauge theories of this type. In our case the presence of the term proportional to a_0^2 in the Lagrangian provides a way to spontaneously break parity and time-reversal symmetry, since this term breaks both gauge and Lorentz symmetries.

We start our analysis by deriving a purely magnetic action. To this end we integrate out N fermionic orbital degrees of freedom using an imaginary time formalism and expand the logarithm of the fermion determinant up to quadratic order in the remaining fields. This leads to the following effective action,

$$S_{\rm eff} = \frac{1}{2}\int\frac{d^3p}{(2\pi)^3}\left[\Pi(p)\left(p^2\delta_{\mu\nu} - p_\mu p_\nu\right)a_\mu(p)a_\nu(-p) + \frac{J^2}{g}a_0(p)a_0(-p)\right], \tag{4}$$

where $p = (\omega, v_F\mathbf{p})$, the Greek indices run from 0 to 2, and $\Pi(p) = NJ^2/(16|p|)$ is the vacuum polarization. Here $|p| = \sqrt{\omega^2 + v_F^2\mathbf{p}^2}$. From Eq. (4) we derive the propagator,

$$D_{\mu\nu}(p) = \langle a_\mu(p)a_\nu(-p)\rangle = \frac{1}{p^2\Pi(p)}\left\{\delta_{\mu\nu} + \left[\frac{g}{J^2}p^2\Pi(p) + 1\right]\frac{p_\mu p_\nu}{\omega^2} - \frac{(p_\mu\delta_{\nu 0} + p_\nu\delta_{\mu 0})}{\omega}\right\}. \tag{5}$$

Interestingly, this result shows that a vector field with a mass term only along the temporal direction is not gapped. With an isotropic mass term of the form $(J^2/g)a_\mu a_\mu$, we would obtain instead $D_{\mu\nu}(p) = [p^2\Pi(p) + J^2/g]^{-1}[\delta_{\mu\nu} + (g/J^2)\Pi(p)p_\mu p_\nu]$, which is clearly gapped. As we will see shortly, this difference is important, as in our case two of the components of the vector field relate to the magnetization and magnetic excitations are supposed to be gapless.

The propagator (5) does not smoothly connect to the strongly coupled regime, $g \to \infty$. This is a typical behavior for massive vector fields [21] which is also reflected here, although our vector field is only massive along the temporal direction. Note, however, that in the strongly coupled regime our model reduces to a QED model in 2+1 dimensions with two-component Dirac fermions. As mentioned earlier, no CS term is generated in this case when the Dirac fermions become gapped [39].

The purely magnetic effective action is finally obtained by integrating out a_0 in the effective action (4). This yields,

$$\begin{aligned} S_{\rm eff}^{\rm FM} &= \frac{1}{2}\int \frac{d^3p}{(2\pi)^3}\Pi(p)\left[p^2\delta_{ij} - F(p)v_F^2 p_i p_j\right]a_i(p)a_j(-p) \\ &= \frac{1}{2}\int \frac{d^3p}{(2\pi)^3}\Pi(p)\left\{\left[p^2 - F(p)v_F^2\mathbf{p}^2\right]\mathbf{n}(p)\cdot\mathbf{n}(-p)\right. \\ &\quad \left. + \, F(p)v_F^2[\mathbf{p}\cdot\mathbf{n}(p)][\mathbf{p}\cdot\mathbf{n}(-p)]\right\}, \end{aligned} \tag{6}$$

where

$$F(p) = \frac{p^2\Pi(p) + J^2/g}{v_F^2\mathbf{p}^2\Pi(p) + J^2/g}. \tag{7}$$

Going back to real time, the magnetic susceptibility $\chi(\omega, \mathbf{p}) = \langle n_+(\omega, \mathbf{p})n_-(\omega, \mathbf{p})\rangle$, where $n_\pm = n_x \pm n_y$, is determined from Eq. (6) as,

$$\begin{aligned} \chi(\omega, \mathbf{p}) &= \frac{16}{NJ^2\sqrt{v_F^2\mathbf{p}^2 - (\omega + i\delta)^2}} \\ &\times \left\{1 - \frac{Ngv_F^2\mathbf{p}^2}{(\omega + i\delta)^2}\left[1 + \frac{16}{Ng}\frac{\sqrt{v_F^2\mathbf{p}^2 - (\omega + i\delta)^2}}{v_F^2\mathbf{p}^2}\right]\right\} \end{aligned} \tag{8}$$

From the pole of $\chi(\omega, \mathbf{p})$ we infer that the spin-wave velocity is identical to the Fermi velocity. Furthermore, by comparing with the scaling behavior $\chi(\omega, \mathbf{p}) \sim [v_F^2\mathbf{p}^2 - (\omega + i\delta)^2]^{\eta/2-1}$ for ω near $v_F|\mathbf{p}|$, yields an anomalous scaling dimension $\eta = 1$. This induced anomalous dimension on the topological surface is very different from the one of a two-dimensional planar FM at $T = 0$ corresponding to a three-dimensional ($d = 2 + 1$) XY universality class, having $\eta \approx 0.04$.

Next, we consider the fermionic propagator. Within an imaginary time formalism, the fermion propagator $G(p)$ is given in general form by

$$G^{-1}(p) = i\gamma_\mu p_\mu + J\int \frac{d^3k}{(2\pi)^3}\gamma_\mu G(p - k)D_{\mu\nu}(k)\Gamma_\nu(p, k), \tag{9}$$

where $\Gamma_\nu(p, k)$ is the vertex function. It is understood that the Dirac matrices above are the imaginary time counterparts of the real time ones defined earlier. They are assumed to satisfy the Clifford algebra $\gamma_\mu\gamma_\nu + \gamma_\nu\gamma_\mu = 2\delta_{\mu\nu}$. In order to determine $G(p)$ approximately, we make the decomposition $G^{-1}(p) = Z(p)i\gamma_\mu p_\mu + \Sigma(p)$ and assume the lowest order form for the vertex function, $\Gamma_\mu(p, k) = J\gamma_\mu$. Furthermore, we will set $Z(p) \approx 1$ for $G(p)$ inside the integral in Eq. (9). A mass will be generated if $m \equiv \Sigma(0)$ does not vanish. Note that a non-vanishing m implies that $\langle\bar\psi\psi\rangle \neq 0$. Since $\bar\psi\psi = n_\uparrow - n_\downarrow$, mass generation implies also an emergent third component of the magnetization. Our strategy will be to make an approximation in which $\Sigma(p)$ is uniform, $\Sigma(p) = \Sigma(0) = m$, and see whether there is a solution to Eq. (9) with $m \neq 0$. We will solve Eq. (9) under the assumption that $m \ll |p| \ll \Lambda$, where $\Lambda \approx NJ^2$ is an ultraviolet cutoff. The fermion mass modifies the vacuum polarization. However, under the assumption $m \ll |p| \ll \Lambda$, the contribution that is even under parity and time-reversal, corresponding to the transverse term in Eq. (4), remains unchanged. On the other hand, an additional term odd under parity is generated. In order to investigate the gap equation, we follow Ref. [39] and assume that $N - L$ fermions acquire a positive mass $+m$, while the remaining L fermions acquire a negative mass $-m$. Thus, the effective action (4) for the vector field receives the following additional contribution odd under parity and time-reversal,

$$S_{\rm eff}^{\rm odd} = \frac{2}{N}\sum_{i=}^{N} m_i \int \frac{d^3p}{(2\pi)^3}\Pi(p)\epsilon_{\mu\nu\lambda}p_\lambda a_\mu(p)a_\nu(-p). \tag{10}$$

This leads in turn to an additional term in the vector field propagator given by $D_{\mu\nu}^{\rm odd}(p) = -32\sum_i(m_i/N)\epsilon_{\mu\nu\lambda}p_\lambda/(NJ^2|p|^3)$. Thus, the following self-consistent equation for m_i is obtained,

$$\begin{aligned} 1 = \frac{16}{N}&\left\{\int \frac{d^3k}{(2\pi)^3}\frac{1}{|k|(k^2 + m^2)} + \int \frac{d^3k}{(2\pi)^3}\frac{|k|}{\omega^2(k^2 + m^2)}\right. \\ &\left. - \, 2\,\frac{m}{m_i}\left(\frac{N - 2L}{N}\right)\int \frac{d^3k}{(2\pi)^3}\frac{1}{|k|(k^2 + m^2)}\right\} \\ &+ g\int \frac{d^3k}{(2\pi)^3}\frac{k^2}{\omega^2(k^2 + m^2)}. \end{aligned} \tag{11}$$

The integral in ω in the last term of Eq. (11) needs to be regularized due to the pole at $\omega = 0$. To this end, we add a small imaginary part $\omega \to \omega + i\delta$ and let $\delta \to 0$ at the end.

After performing all integrals, the gap equation for $i = 1, \ldots, N - L$ becomes,

$$1 = \frac{16}{\pi^2 N}\left(1 - \frac{2L}{N}\right)\ln(m/\Lambda) + \frac{g\Lambda}{4\pi}\left(1 + \frac{1}{\pi} - 3\frac{|m|}{\Lambda}\right), \tag{12}$$

while for $i = N - L + 1, \ldots, N$, we obtain,

$$1 = -\frac{16}{\pi^2 N}\left(1 - \frac{2L}{N}\right)\ln(m/\Lambda) + \frac{g\Lambda}{4\pi}\left(1 + \frac{1}{\pi} - 3\frac{|m|}{\Lambda}\right). \tag{13}$$

These gap equations are only compatible with each other if N is even, i.e., $L = N/2$. If we introduce the dimensionless quantities $\hat m = m/\Lambda$ and $\hat g = \Lambda g$, we obtain for an even number of Dirac fermions,

$$|\hat m| = \frac{\pi + 1}{3\pi}\left(1 - \frac{\hat g_c}{\hat g}\right), \tag{14}$$

where

$$\hat g_c = \frac{4\pi^2}{\pi + 1}\left(1 - \frac{8\ln 2}{\pi^2 N}\right). \tag{15}$$

Thus, we obtain that if N is even a gap is generated provided $\hat{g} > \hat{g}_c$. In this case there is no generation of CS term. Therefore, parity and time-reversal symmetries remain preserved. The above result distinguishes itself from the QED case [39] due to a complete cancellation of the logarithm.

The above analysis does not necessarily imply that there is no mass generation for an odd number of Dirac fermions. Actually, *for all N* there is in principle the possibility that $m_i = +m$ for all i. In this case the logarithmic term survives and dominates for $|m| \ll \Lambda$ over the linear term in $|m|$. Thus, the gap equation is given by Eq. (12) with $L = 0$ and where the term proportional to $|m|/\Lambda$ is neglected. Therefore, we obtain,

$$\hat{m} = \exp\left[-\frac{(\pi+1)N}{64}(\hat{g} - \hat{g}_c)\right], \tag{16}$$

which is only meaningful for $\hat{g} > \hat{g}_c$, otherwise it does not decrease with increasing N, which would at large N contradict the condition $m \ll |p| \ll \Lambda$, in a situation reminiscent from the QED case [39]. However, in our case it is possible to overcome this difficulty and obtain in addition the generation of a CS term. Although Eq. (16) is in principle valid for any N, including the even ones, the solution (14) is energetically more stable for N even, as it still gives a nonzero gap even in the strong-coupling regime $\hat{g} \to \infty$.

In conclusion, we have shown that in topological insulators with a proximity-induced in-plane magnetization a gap can be spontaneously generated by tuning the local electronic interaction above a critical value, leading in this way to a semimetal-insulator transition. We find that when the number of Dirac fermions N is even, the masses are generated in pairs $\pm|m|$ and no CS term is generated, so that parity is overall preserved. On the other hand, for N odd all generated masses are equal and positive, so that a CS term is generated. That no CS term is generated for N even is physically reasonable, since in this case we can change to a representation where there are $N/2$ four-component Dirac spinors, in which case the model may be reinterpreted as some model for graphene, a material featuring an even number of Dirac cones. Interestingly, in such a graphene model the gap generation is associated to a mass spectrum containing masses $\pm|m|$, a scenario not considered so far in interacting models for graphene where the vector field has only the time component [22, 23]. TIs, on the other hand, have an odd number of Dirac cones. In this context, recent experiments on EuS/Bi$_2$Se$_3$ heterostructures open the possibility that the experimentally elusive gap generation in QED-like theories may finally be observed in the near future.

We would like to thank Igor Herbut, Jörg Schmalian, Peng Wei, and Jagadeesh Moodera for interesting discussions. We thank J. S. Moodera for sending us his preprint prior publication. The authors would like to thank the Deutsche Forschungsgemeinschaft (DFG) for the financial support via the SFB TR 12.

---

[1] M. Z. Hasan and C.L. Kane, Rev. Mod. Phys. **82**, 3045 (2010).
[2] X.-L. Qi and S.-C. Zhang, Rev. Mod. Phys. **83**, 1057 (2011).
[3] D. Pesin and A. H. MacDonald, Nature Mat. **11**, 409 (2012).
[4] X.-L. Qi, T. L. Hughes, and S.-C. Zhang, Phys. Rev. B **78**, 195424 (2008).
[5] T. Yokoyama, J. Zang, and N. Nagaosa, Phys. Rev. B **81**, 241410(R) (2010).
[6] K. Nomura and N. Nagaosa, Phys. Rev. B **82**, 161401(R) (2010).
[7] I. Garate and M. Franz, Phys. Rev. Lett. **104**, 146802 (2010).
[8] G. Rosenberg and M. Franz, Phys. Rev. B **82**, 035105 (2010).
[9] C. Wickles and W. Belzig, Phys. Rev. B **86**, 035151 (2012).
[10] Y. Tserkovnyak and D. Loss, Phys. Rev. Lett. **108**, 187201 (2012).
[11] F. S. Nogueira and I. Eremin, Phys. Rev. Lett. **109**, 237203 (2012).
[12] L. Oroszlány and A. Cortijo, Phys. Rev. B **86**, 195427 (2012).
[13] W. Luo and X.-L. Qi, Phys. Rev. B **87**, 085431 (2013).
[14] Y. S. Hor, P. Roushan, H. Beidenkopf, J. Seo, D. Qu, J. G. Checkelsky, L. A. Wray, D. Hsieh, Y. Xia, S.-Y. Xu, D. Qian,, M. Z. Hasan, N. P. Ong, A. Yazdani, and R. J. Cava, Phys. Rev. B **81**, 195203 2010.
[15] L. A. Wray , S.-Y. Xu , Y. Xia , D. Hsieh , A. V. Fedorov , Y. S. Hor , R. J. Cava , A. Bansil , H. Lin, and M. Z. Hasan, Nature Phys. **7**, 32 (2011).
[16] I. Vobornik, U. Manju, J. Fujii, F. Borgatti, P. Torelli, D. Krizmancic, Y. S. Hor, R. J. Cava, and G. Panaccione, Nano Lett. **11**, 4079 (2011).
[17] M. R. Scholz, J. Sánchez-Barriga, D. Marchenko, A. Varykhalov, A. Volykhov, L. V. Yashina, and O. Rader, Phys. Rev. Lett. **108**, 256810 (2012).
[18] J. G. Checkelsky, J. Ye , Y. Onose, Y. Iwasa1, and Y. Tokura, Nat. Phys. **8**, 729 (2012).
[19] P. P. J. Haazen, J.-B. Laloë, T. J. Nummy, H. J. M. Swagten, P. Jarillo-Herrero, D. Heiman, and J. S. Moodera, Appl. Phys. Lett. **100**, 082404 (2012).
[20] P. Wei, F. Katmis, B. A. Assaf, H. Steinberg, P. Jarillo-Herrero, D. Heiman, and J. S. Moodera, to appear in the Phys. Rev. Lett. (2013).
[21] See Chapter 18 in: J. Zinn-Justin, *Quantum Field Theory and Critical Phenomena*, 2nd ed. (Oxford University Press, Oxford, 1993).
[22] I. F. Herbut, Phys. Rev. Lett. **97**, 146401 (2006).
[23] D. E. Sheehy and J. Schmalian, Phys. Rev. Lett. **99**, 226803 (2007).
[24] I. F. Herbut, V. Juricić, and B. Roy, Phys. Rev. B **79**, 085116 (2009).
[25] W. Thirring, Ann. Phys. (N.Y.) **3**, 91 (1958).
[26] D. H. Kim and P. A. Lee, Ann. Phys. (N.Y.) **272**, 130 (1999).
[27] W. Rantner and W.-G. Wen, Phys. Rev. Lett. **86**, 3871 (2001); Phys. Rev. B **66**, 144501 (2002).
[28] Z. Tesanovic, O. Vafek, and M. Franz, Phys. Rev. B **65**, 180511 (2002).
[29] I. F. Herbut, Phys. Rev. Lett. **88**, 047006 (2002); Phys. Rev. B **66**, 094504 (2002).
[30] D. V. Khveschchenko, Phys. Rev. Lett. **87**, 246802 (2001).
[31] V. P. Gusynin, V. A. Miransky, S. G. Shaparov and I. A. Shovkovy, Phys. Rev. B **74**, 195429 (2006).
[32] J. A. Drut and T. A. Lähde, ; Phys. Rev. Lett. **102**, 026802 (2009); Phys. Rev. B **79**, 165425 (2009).
[33] Z. Y. Meng, T. C. Lang, S. Wessel, F. F. Assaad, and A. Mura-

matsu, Nature **464**, 847 (2010).

[34] P. A. Lee., N. Nagaosa, and X.-G. Wen, Rev. Mod. Phys. **78**,17 (2006).

[35] M. Franz and Z. Tesanovic, Phys. Rev. Lett. **87**, 257003 (2001); M. Franz, Z. Tesanovic, and O. Vafek, Phys. Rev. B **66**, 054535 (2002).

[36] T. Appelquist and L.C.R. Wijewardhana, Phys. Rev. Lett. **60**, 2575 (1988); D. Nash, Phys. Rev. Lett. **62**, 3024 (1989); K.-I. Kondo and P. Maris, Phys. Rev. D **52**, 1212 (1995); P. Maris, Phys. Rev. D **54**, 4049 (1996); W. Armour, J. B. Kogut, and C. Strouthos, Phys. Rev. D **82**, 014503 (2010).

[37] R.D. Pisarski, Phys. Rev. D **29**, R2423 (1984); T. W. Appelquist, M. Bowick, D. Karabali, and L. C. R. Wijewardhana, Phys. Rev. D **33**, 3704 (1986).

[38] S. Deser, R. Jackiw, and S. Templeton, Phys. Rev. Lett. **48**, 975 (1982); A. N. Redlich, Phys. Rev. D **29**, 2366 (1984).

[39] T. Appelquist, M.J. Bowick, D. Karabali, and L.C.R. Wijewardhana, Phys. Rev. D **33**, 3774 (1986).

[40] P. M. Lo and E. S. Swanson, Phys. Rev. D **83**, 065006 (2011).

[41] S.R. Coleman and B.R. Hill, Phys. Lett. B **159**, 184 (1985).

[42] C. Vafa and E. Witten, Commun. Math. Phys. **95**, 257 (1984); Nucl. Phys. B **234**, 173 (1984); this result was revisited some time ago by M. B. Einhorn and J. Wudka, Phys. Rev. D **67**, 045004 (2003).